

## **Appendix 19**

### Site Synopses

## Appendix 20 Site Synopses

### Great Island Channel SAC Site Code: 001058

Site Name: Great Island Channel SAC Site Code: 001058 The Great Island Channel stretches from Little Island to Midleton, with its southern boundary being formed by Great Island. It is an integral part of Cork Harbour which contains several other sites of conservation interest. Geologically, Cork Harbour consists of two large areas of open water in a limestone basin, separated from each other and the open sea by ridges of Old Red Sandstone. Within this system, Great Island Channel forms the eastern stretch of the river basin and, compared to the rest of Cork Harbour, is relatively undisturbed. Within the site is the estuary of the Owennacurra and Dungourney Rivers. These rivers, which flow through Midleton, provide the main source of freshwater to the North Channel. The site is a Special Area of Conservation (SAC) selected for the following habitats and/or species listed on Annex I / II of the E.U. Habitats Directive (\* = priority; numbers in brackets are Natura 2000 codes):

[1140] Tidal Mudflats and Sandflats

[1330] Atlantic Salt Meadows

The main habitats of conservation interest in Great Island Channel SAC are the sheltered tidal sand and mudflats and the Atlantic salt meadows. Owing to the sheltered conditions, the intertidal flats are composed mainly of soft muds. These muds support a range of macro-invertebrates, notably *Macoma balthica*, *Scrobicularia plana*, *Hydrobia ulvae*, *Nephtys hombergi*, *Nereis diversicolor* and *Corophium volutator*. Green algal species occur on the flats, especially *Ulva lactuca* and *Enteromorpha* spp. Cordgrass (*Spartina* spp.) has colonised the intertidal flats in places, especially at Rossleague and Belvelly. The saltmarshes are scattered through the site and are all of the estuarine type on mud substrate. Species present include Sea Purslane (*Halimione portulacoides*), Sea Aster (*Aster tripolium*), Thrift (*Armeria maritima*), Common Saltmarsh-grass (*Puccinellia maritima*), Sea Plantain (*Plantago maritima*), Greater Sea-spurrey (*Spergularia media*), Lax-flowered Sea-lavender (*Limonium humile*), Sea Arrowgrass (*Triglochin maritimum*), Sea Mayweed (*Matricaria maritima*) and Red Fescue (*Festuca rubra*).

The site is extremely important for wintering waterfowl and is considered to contain three of the top five areas within Cork Harbour, namely North Channel, Harper's Island and Belvelly-Marino Point. Shelduck is the most frequent duck species with 800-1,000 birds centred on the Fota/Marino Point area. There are also large flocks of Teal and Wigeon, especially at the eastern end. Waders occur in the greatest density north of Rosslare, with Dunlin, Godwit, Curlew and Golden Plover the commonest species. A population of about 80 Grey Plover is a notable feature of the area. All the mudflats support feeding birds; the main roost sites are at Weir Island and Brown Island, and to the north of Fota at Killacloyne and Harper's Island. Ahanesk supports a roost also but is subject to disturbance. The numbers of Grey Plover and Shelduck, as given above, are of national importance. The site is an integral part of Cork Harbour which is a wetland of international importance for the birds it supports.

Overall, Cork Harbour regularly holds over 20,000 waterfowl and contains internationally important numbers of Black-tailed Godwit (1,181) and Redshank (1,896), along with nationally important numbers of nineteen other species. Furthermore, it contains large Dunlin (12,019) and Lapwing (12,528) flocks. All counts are average peaks, 1994/95 – 1996/97. Much of the site falls within Cork Harbour Special Protection Area, an important bird area designated under the E.U. Birds Directive. While the main land use within the site is aquaculture (oyster farming), the greatest threats to its conservation significance come from road works, infilling, sewage outflows and possible marina developments. The site is of major importance for the two habitats listed on Annex I of the E.U.

Habitats Directive, as well as for its important numbers of wintering waders and wildfowl. It also supports a good invertebrate fauna.

Version date: 24.09.2013 2 of 2 001058\_Rev13.Doc

**SITE NAME: CORK HARBOUR SPA SITE CODE: 004030**

Cork Harbour is a large, sheltered bay system, with several river estuaries - principally those of the Rivers Lee, Douglas, Owenboy and Owennacurra. The SPA site comprises most of the main intertidal areas of Cork Harbour, including all of the North Channel, the Douglas River Estuary, inner Lough Mahon, Monkstown Creek, Lough Beg, the Owenboy River Estuary, Whitegate Bay, Ringabella Creek and the Rostellan and Poul nabibe inlets. Owing to the sheltered conditions, the intertidal flats are often muddy in character. These muds support a range of macro-invertebrates, notably *Macoma balthica*, *Scrobicularia plana*, *Hydrobia ulvae*, *Nephtys hombergi*, *Nereis diversicolor* and *Corophium volutator*. Green algae species occur on the flats, especially *Ulva* spp. Cordgrass (*Spartina* spp.) has colonised the intertidal flats in places, especially where good shelter exists, such as at Rossleague and Belvelly in the North Channel.

Salt marshes are scattered through the site and these provide high tide roosts for the birds. Some shallow bay water is included in the site. Rostellan Lake is a small brackish lake that is used by swans throughout the winter. The site also includes some marginal wet grassland areas used by feeding and roosting birds. The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for the following species: Little Grebe, Great Crested Grebe, Cormorant, Grey Heron, Shelduck, Wigeon, Teal, Mallard, Pintail, Shoveler, Redbreasted Merganser, Oystercatcher, Golden Plover, Grey Plover, Lapwing, Dunlin, Black-tailed Godwit, Bar-tailed Godwit, Curlew, Redshank, Greenshank, Blackheaded Gull, Common Gull, Lesser Black-backed Gull and Common Tern. The site is also of special conservation interest for holding an assemblage of over 20,000 wintering waterbirds.

The E.U. Birds Directive pays particular attention to wetlands and, as these form part of this SPA, the site and its associated waterbirds are of special conservation interest for Wetland & Waterbirds. Cork Harbour is an internationally important wetland site, regularly supporting in excess of 20,000 wintering waterfowl. Of particular note is that the site supports internationally important populations of Black-tailed Godwit (1,896) and Redshank (2,149) - all figures given are five year mean peaks for the period 1995/96 to 1999/2000. Nationally important populations of the following 19 species occur: Little Grebe (57), Great Crested Grebe (253), Cormorant (521), Grey Heron (80), Shelduck (2,009), Wigeon (1,791), Teal (1,065), Mallard (513), Pintail (57), Shoveler (103), Red-breasted Merganser (121), Oystercatcher (1,809), Golden Plover (3,342), Grey Plover (95), Lapwing (7,569), Dunlin (9,621), Bartailed Godwit (233), Curlew (2,237) and Greenshank (46). The Shelduck population is the largest in the country (over 10% of national total). Other species using the site include Mute Swan (38), Whooper Swan (5), Pochard (72), Gadwall (6), Tufted Duck (64), Goldeneye (21), Coot (53), Ringed Plover (73), Knot (26) and Turnstone (113). Cork Harbour is an important site for gulls in winter and autumn, especially Black-headed Gull (3,640), Common Gull (1,562) and Lesser Black-backed Gull (783), all of which occur in numbers of national importance.

Little Egret and Mediterranean Gull, two species which have recently colonised Ireland, also occur at this site. A range of passage waders occurs regularly in autumn, including such species as Ruff (5-10), Spotted Redshank (1-5) and Green Sandpiper (1-5). Numbers vary between years and usually a few of each of these species over-winter. Cork Harbour has a nationally important breeding colony of

Common Tern (102 pairs in 1995). The birds have nested in Cork Harbour since about 1970, and since 1983 on various artificial structures, notably derelict steel barges and the roof of a Martello Tower. The birds are monitored annually and the chicks are ringed. Cork Harbour is of major ornithological significance, being of international importance both for the total numbers of wintering birds (i.e. > 20,000) and also for its populations of Black-tailed Godwit and Redshank. In addition, it supports nationally important wintering populations of 22 species, as well as a nationally important breeding colony of Common Tern. Several of the species which occur regularly are listed on Annex I of the E.U. Birds Directive, i.e. Whooper Swan, Little Egret, Golden Plover, Bar-tailed Godwit, Ruff, Mediterranean Gull and Common Tern. The site provides both feeding and roosting sites for the various bird species that use it. Cork Harbour is also a Ramsar Convention site and part of Cork Harbour SPA is a Wildfowl Sanctuary.

21.1.2015

#### **Ballycotton Bay SPA Site Code: 004022**

Situated on the south coast of Co. Cork, Ballycotton Bay is an east-facing coastal complex, which stretches northwards from Ballycotton to Ballynamona, a distance of c. 2 km. The site comprises two sheltered inlets which receive the flows of several small rivers. The southern inlet had formerly been lagoonal (Ballycotton Lake) but breaching of the shingle barrier in recent times has resulted in the area reverting to an estuarine system. The principal habitat within the site is inter-tidal sand and mudflats. These are mostly well-exposed and the sediments are predominantly firm sands. In the more sheltered conditions of the inlets, sediments contain a higher silt fraction. The inter-tidal flats provide the main feeding habitat for the wintering birds. Sandy beaches are well represented. Salt marshes fringe the flats in the sheltered inlets and these provide high tides roosts. A small area of shallow marine water is also included. The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for the following species: Teal, Ringed Plover, Golden Plover, Grey Plover, Lapwing, Black-tailed Godwit, Bar-tailed Godwit, Curlew, Turnstone, Common Gull and Lesser Black-backed Gull. The E.U. Birds Directive pays particular attention to wetlands, and as these form part of this SPA, the site and its associated waterbirds are of special conservation interest for Wetland & Waterbirds.

Ballycotton Bay supports an excellent diversity of wintering waterbird species. The site supports nationally important populations of the following species: Teal (903), Ringed Plover (167), Golden Plover (2,383), Grey Plover (124), Lapwing (2,782), Black-tailed Godwit (136), Bar-tailed Godwit (175), Curlew (853), Turnstone (179), Common Gull (584) and Lesser Black-backed Gull (1,293) - all figures are five year mean peaks for the period 1995/96 to 1999/2000. Other species which occur include Shelduck (99), Wigeon (522), Mallard (232), Oystercatcher (255), Dunlin (575), Sanderling (56), Redshank (117), Greenshank (12) and Great Black-backed Gull (324). Ballycotton Bay was formerly utilised by Bewick's Swan but the birds have abandoned the site since the reversion of the lagoonal habitat to estuarine conditions. The site is a well-known location for passage waders, especially in autumn. Species such as Ruff, Little Stint, Curlew Sandpiper, Green Sandpiper and Spotted Redshank occur annually though in variable numbers. Small numbers of Ruff may also be seen in late winter and spring. Rarer waders, such as Wood Sandpiper and Pectoral Sandpiper, have also been recorded. While relatively small in area, Ballycotton Bay supports an excellent diversity of wintering waterbirds and has nationally important populations of eleven species, of which two, Golden Plover and Bar-tailed Godwit, are listed on Annex I of the E.U. Birds Directive. Ballycotton Bay is also a Ramsar Convention site and part of the Ballycotton Bay SPA is a Wildfowl Sanctuary.

30.5.2015

### **Sovereign Islands SPA 004124**

The Sovereign Islands are two very small marine islands located approximately 1 km off the coastline at the entrance to Oysterhaven Bay in Co. Cork. The islands are rocky stacks separated by a narrow sound of about 20 m width. The eastern island is flat-topped and rises to 24 m above sea level; the western one is more peaked and rises to 30 m. The geology is Lower Carboniferous limestones and shales. Both islands are largely devoid of soil apart from small amounts of organic matter trapped in cracks. Vegetation is sparse, with plants such as Sea Beet (*Beta vulgaris*), Spurrey (*Spergularia* spp.) and Orache (*Atriplex* spp.) recorded. The surrounding sea, to a distance of 200 m, is included. The site is a Special Protection Area (SPA) under the E.U. Birds Directive, of special conservation interest for the following species: Cormorant.

The islands are important for breeding seabirds, with most occurring on the eastern stack. A Cormorant colony has been known since the late 1960s and 156 pairs were recorded here in 1999. A more recent survey in 2008 recorded 89 pairs. Herring Gull and Great Black-backed Gull also breed, with 10 and 75 pairs respectively in 1999. Sovereign Islands SPA is of ornithological importance mainly for the breeding colony of Cormorant, which is both the largest in Co. Cork and of national importance. The non-migratory population of Great Black-backed Gull is also of national importance.

14.12.2011